Strong Lensing Legacy Survey

Automated search for strong lenses in CFHTLS data

IAP: C. Alard, B. Fort, Y. Mellier, JF. Sygnet

Toulouse: R. Cabanac (Tarbes), G. Soucail

Marseille: JP. Kneib



How to find highly distorded (arc-like) images, using automated softwares ?









Deflection of light by a point mass



General relativity at first order $\theta = \frac{GM}{b}$

Geometry of gravitational lensing



$$\beta = \theta - \frac{D_{ls}}{D_s} \alpha$$
 Lens equation

Point mass lens





Spatially extended lens



Elliptical lens with isothermal profile



Peter Newbury 1997





Finding arcs or arclets in wide field CFHTLS images



Most of the time

somewhat unusual



Modeling arcs



local elongation and orientation at each point of the image

Estimating local geometry



Parameters of local geometry

Estimation of θ not very noisy using 2^{nd} order moment

Variance on θ proportional to noise variance

$$\sigma_{\theta} \simeq \frac{1}{4} \frac{S}{N} \simeq 1^{\circ} \text{ for } \frac{S}{N} = 10$$

Elongation a:/b --> divergent, very sensitive to noise and to details of the local distribution

Estimating local elongation



Marginal Distributions projected along axis

Elongation estimator

$$Q(x_{0}, y_{0}) = \frac{1}{2M} \frac{I_{Y}(x_{0})}{SUP.[I_{X}(x_{0}+x)]_{[-M < x < M]}}$$

Decomposable locally $I(x+x_0, y+y_0) = f(x)g(y)$

$$Q(x_0, y_0) \leq \frac{g(0)}{\int g(y) dy} \qquad g(y) = \alpha \ G(\frac{y}{b}) \qquad Q(x_0, y_0) \leq \frac{G(0)}{b}$$

Bounded estimator, maximal for small b (PSF size) Equal to sup bound if distribution is flat along main axis ~ Arc

Other propriety of the estimator

Homothetic profiles:
$$f(x) = \alpha F(\frac{x}{a})$$
; $g(x) = \beta G(\frac{x}{b})$

$$(a,b) \ll M$$
, $Q(x_0, y_0) = \frac{G(0)}{2M} \frac{a}{b}$

$$a \gg M$$
, $Q(x_0, y_0) \rightarrow \frac{G(0)}{b}$ Saturation value

Noise on estimation of local elongation

Truncated expansion of Q(x,y) for small errors Combination of errors for the 2 marginale distribution using an order statistics

$$\frac{Q}{N_Q} \simeq 0.45 \frac{S}{N}$$

For $\frac{S}{N} = 10$ noise is only ~ 20% of signal Dynamic on Q(x,y) ~ factor 3, noise is not an issue

Illustration with CFHTLS wide image





Small arcs – observational limits





Galaxy groups



Mass ~ 10^{13} Solar mass

Velocity dispersion inside group a few hundreds km/s

Lensing du to galaxy groups, separations of a few arc seconds







Arcs appears near critical radius size of system is close to Einstein radius ----> Estimation of Einstein radius ----> Mass of deflector



Effect of sub-structures, small perturbations of caustics --> arc shape modified col dark matter granularity, visible in arcs ?



Uneasy to disentangle from intrisic source granularity Future

Data acquisition & data reduction

Spectro & redshifts: ESO Proposals, Gemini

Automated search in new CFTHLS releases

Comparison of the observational search with cosmological models – Ray tracing in numerical simulations, reconstruction of arcs, computation of arcs properties & estimation of detection efficiency